

Safe food and feed through an integrated toolbox for mycotoxin management: the MyToolBox approach

R. Krska^{1*}, M. de Nijs², O. McNERNEY³, M. Pichler⁴, J. Gilbert⁵, S. Edwards⁶, M. Suman⁷, N. Magan⁸, V. Rossi⁹, H.J. van der Fels-Klerx², F. Bagi¹⁰, B. Poschmaier¹, M. Sulyok¹, F. Berthiller¹ and H.P. van Egmond¹¹

¹University of Natural Resources and Life Sciences, Vienna (BOKU), Department of Agrobiotechnology (IFA-Tulln), Center for Analytical Chemistry, Konrad-Lorenz-Str. 20, 3430 Tulln, Austria; ²RIKILT Wageningen University & Research, Akkermaalsbos 2, 6708 WB Wageningen, the Netherlands; ³Innovacio i Recerca Industrial i Sostenible SL, Avda. Carl Friedrich Gauss 11, 08860 Castelldefels, Spain; ⁴International Association for Cereal Science and Technology, Marxergasse 2, 1030 Vienna, Austria; ⁵FoodLife International Ltd., ODTU Teknokent, Ikizler Binasi No Ara-1 ODTU, Cankaya 06800, Turkey; ⁶Harper Adams University, Department of Crop and Environment Science, Edgmond, Newport, Shropshire TF10 8NB, United Kingdom; ⁷BARILLA S.p.A., Advanced Laboratory Research, Barilla Research Labs, via Mantova 166, 43122 Parma, Italy; ⁸Cranfield University, Applied Mycology Group, Soil and Agrifood Institute, Cranfield Health, Cranfield, Bedford MK43 0AL, United Kingdom; ⁹HORTA SRL, via E. Gorra, 29122 Piacenza, Italy; ¹⁰University of Novi Sad, Faculty of Agriculture, Trg Dositeja Obradovića 8, Novi Sad 21000, Serbia; ¹¹Retired from RIKILT, Wageningen University & Research, the Netherlands; rudolf.krska@boku.ac.at

Received: 13 September 2016 / Accepted: 4 October 2016

© 2016 Wageningen Academic Publishers

OPEN ACCESS



OPINION ARTICLE

Abstract

There is a pressing need to mobilise the wealth of knowledge from the international mycotoxin research conducted over the past 25-30 years, and to perform cutting-edge research where knowledge gaps still exist. This knowledge needs to be integrated into affordable and practical tools for farmers and food processors along the chain in order to reduce the risk of mycotoxin contamination of crops, feed and food. This is the mission of MyToolBox – a four-year project which has received funding from the European Commission. It mobilises a multi-actor partnership (academia, farmers, technology small and medium sized enterprises, food industry and policy stakeholders) to develop novel interventions aimed at achieving a significant reduction in crop losses due to mycotoxin contamination. Besides a field-to-fork approach, MyToolBox also considers safe use options of contaminated batches, such as the efficient production of biofuels. Compared to previous efforts of mycotoxin reduction strategies, the distinguishing feature of MyToolBox is to provide the recommended measures to the end users along the food and feed chain in a web-based MyToolBox platform (e-toolbox). The project focuses on small grain cereals, maize, peanuts and dried figs, applicable to agricultural conditions in the EU and China. Crop losses using existing practices are being compared with crop losses after novel pre-harvest interventions including investigation of genetic resistance to fungal infection, cultural control (e.g. minimum tillage or crop debris treatment), the use of novel biopesticides suitable for organic farming, competitive biocontrol treatment and development of novel modelling approaches to predict mycotoxin contamination. Research into post-harvest measures includes real-time monitoring during storage, innovative sorting of crops using vision-technology, novel milling technology and studying the effects of baking on mycotoxins at an industrial scale.

Keywords: pre-harvest and post-harvest management, integrated mycotoxin management, thermal processing, decision support, dried distillers' grain soluble

1. Introduction

The occurrence of fungal and subsequently, mycotoxin contamination in various crops is of major concern since it has significant implications for food and feed safety, food security and international trade. The European Commission pointed out that 5-10% of global crop production is lost annually due to mycotoxin contamination (EC, 2015). Considering that the European Union (EU) produces about 133 million tonnes (MT) of wheat (~29,038 M€), 68 MT maize (~13,571 M€) and 8 MT oats (~1,543 M€) annually (Eurostat, 2015a,b), about 1.2-2.4 billion Euro can be lost for wheat alone, thus a reduction in these losses of only 1% could save 12-24 M€. The total costs of mycotoxin contamination due to reduced yields, food and feed losses, increased costs for inspection and analyses, and others, may easily reach billions of Euros annually as estimated by Wu and Munkvold (2008). Moreover, with 'one in eight persons worldwide suffering from chronic undernourishment' (Schmale and Munkvold, 2009) and recent reports of the impact of aflatoxins on stunting of children in Africa (Matacic, 2016; Wild *et al.*, 2015), there is clearly a moral obligation to curb these significant food and feed losses.

In a recent multi-mycotoxin survey, 81% of all samples were contaminated with at least one mycotoxin and 45% contained more than one secondary metabolite of fungi (Nährer and Kovalsky, 2014). The EU's Rapid Alert System for Food and Feed (RASFF) (EC, 2016) showed that of the total border rejections in 2015, 18.3% were due to mycotoxin contamination exceeding the EU legislative limits, accounting for the most frequently reported chemical hazard. The product category with the highest susceptibility of aflatoxin contamination was 'nuts, nut products and seeds'. For example, over the last five years RASFF reported 308 alerts relating to aflatoxin levels exceeding legal limits in peanuts imported from China. However, as illustrated by

the RASFF notifications (Figure 1), a significant reduction of about 50% of mycotoxin-related RASFF notifications has been observed over the last 12 years, from 876 notifications in 2004 to 492 notifications in 2015.

Nonetheless, prevention and control of these toxic secondary metabolites remains difficult and the agriculture and food industries continue to be vulnerable to problems of contamination. For example, from February to March 2013, Romania, Serbia and Croatia reported high aflatoxin M₁ contamination of milk. Severe droughts in Serbia in 2012 resulted in 70% of the maize crop being contaminated with aflatoxins (Anonymous, 2013). Feeding this maize to dairy cattle led to high levels of aflatoxin M₁ in milk, up to twice the EU legal limit of 0.05 µg/kg. The milk scandal was fuelled when the permitted level of aflatoxin M₁ in milk in Serbia was temporarily raised to 0.5 µg/kg milk, 10 times the EU legal limits. However, the catastrophic floods and the rainy summer in 2014 resulted in low levels of aflatoxin B₁ with high levels of deoxynivalenol (DON) in Serbia (Jakšić *et al.*, 2015). Obviously, extreme weather conditions as a result of climate change are increasingly affecting the mycotoxin map in Europe and also world-wide (Battilani *et al.*, 2016).

Furthermore, a dramatic increase in DON contamination in grains cultivated in Nordic countries has been observed in recent years. The Norwegian Scientific Committee for Food Safety concluded in a study published in 2013 that '[t]he estimated mean and high (95-percentile) exposures to DON [...] were in the range of, or exceeded the TDI [Tolerable Daily Intake] by almost 2 times in 1-year-old infants and 2-year-old children and up to 3.5 times in years with high DON levels which has been especially true for oat based products (Norwegian Scientific Committee for Food Safety, 2013). In general, grain-based foods (e.g. pasta, bread, bakery products) account for the largest contribution to mycotoxin exposure in all age classes of the EU population,




Figure 1. RASFF mycotoxin notifications from 2004-2015.

in particular due to the mycotoxins produced by *Fusarium* spp. (DON, T-2/HT-2 toxins, zearalenone, ochratoxin A and fumonisins). However, as shown above, contamination with these mycotoxins is difficult to predict and manage, due to new trends in mycotoxin occurrences.

Exposure of livestock to mycotoxins is also estimated to result in substantial economic losses (Wu and Munkvold, 2008). While mycotoxicoses in farmed animals are regularly reported (Jovanović *et al.*, 2015), it is difficult to obtain an accurate estimate of the burden of market loss associated with animal health and productivity, and loss of animal feed. However, several publications from different parts of the world provide some insight into the potential magnitude of the problem. In the USA, Vardon *et al.* (2003) estimated a loss of 225 million US\$ per annum for feed maize due to contamination with aflatoxins, and an additional loss of livestock of 4 million US\$ per annum. Wu and Munkvold (2008) estimated the losses due to decreased weight gain in pigs in the USA, resulting from feeding dried distillers' grain solubles (DDGS) contaminated with fumonisins, at 147 million US\$ annually (assuming 20% DDGS in the diets and market coverage is 100%). A study conducted in Thailand (Lubulwa *et al.*, 2015) revealed that aflatoxins in maize can be expected to cause losses for the livestock sector between 7 and 110 million US\$ per annum. These estimates show that, although limited to aflatoxins and fumonisins in two countries only, the losses in the livestock sector will be significantly multiplied if other commodities and other mycotoxins are taken into account.

Previous success in reducing mycotoxin contamination in food and feed now necessitates the mobilisation of the wealth of knowledge gained over the past 25-30 years, to merge in cutting-edge research and the closure of knowledge gaps. With practical and affordable tools that build on existing research-results, stakeholders along the food and feed chain can reduce the risk of mycotoxin contamination in crops, food and feed. These tools prevent losses and waste along the food and feed chain, and provide traceable information to the supply chain and consumers using mainstream information and communication technology (ICT). Existing knowledge combined with novel findings need to be adapted and integrated to provide the vehicle needed to practically implement this knowledge into tools for use along the food and feed chain. This is the mission of MyToolBox (www.mytoolbox.eu), a four year project, which has been recently funded by the European Commission (EC) and which was officially launched in March 2016 (Figure 2). The project applies a multi-actor and multi-disciplinary approach throughout the food and feed chain with 40% industry participation including five end users and three Chinese institutions who collaborate closely with farmers and other relevant stakeholders from the industry.

2. Objectives

The overall objective of the MyToolBox project is to develop a series of integrated measures, which will enable a significant reduction of different kind of losses due to mycotoxin contamination. Besides a field-to-fork approach along the food and feed chain, MyToolBox also considers safe use options of mycotoxin-contaminated batches such as microbial energy conversion to efficiently produce biogas and bioethanol. The consideration of the entire production chain (soil-field-crop-food processing-waste management-alternative energy-government) is a major motivation behind MyToolBox to ensure food and feed security and safety within a sustainable economic environment. The mycotoxin commodity combinations that are specifically addressed are the most prevalent *Fusarium* mycotoxins (DON, T-2/HT-2 toxins, zearalenone and fumonisins) in wheat, oats, maize and animal feed chains, ochratoxin A in wheat, and aflatoxins in maize, peanuts and dried figs.

The pre- and post-harvest interventions that are being developed within the MyToolBox project for these mycotoxins build on, and integrate, the wealth of knowledge resulting from the substantial EU-, nationally and internationally, currently and previously funded research projects on mycotoxin prevention and control, such as the Mycotoxin Prevention Cluster (FP5 projects), MycoGlobe (FP6 project, MycoRed and MycoKey [www.mycored.eu/d/84/MycoKey/]). Consequently, existing knowledge, novel findings and technologies developed in MyToolBox will be integrated into a comprehensive series of tools and applications. These will be accessible via a dynamic web-based MyToolBox e-platform (MyToolBox; Figure 2) that will also be accessible with mobile devices, which will provide the end users along the whole food and feed chain with the recommended measures for mycotoxin reduction.

By collaborating with partners from industry, farming communities, agronomists, manufacturers and other professionals in agriculture and food processing, SMEs and academia from the EU and beyond, the usability and applicability of the MyToolBox decision support system is being ensured.

The MyToolBox project also anticipates legislative implications of its project outputs. As such, it is cooperating with institutions experienced in standard setting, such as the European Food Safety Authority (EFSA) and the EC's Directorate-General DG Santé. In addition, it aims to develop a sound scientific basis for setting authorisation standards of mycotoxin detoxifying feed additives in China.

The following major objectives are pursued over the course of the 48-month project.




Figure 2. Overview of the MyToolBox project approach (© IRIS).

Pre-harvest objectives

- To reduce fungal infection of grains on field, biopesticides (e.g. Forrer *et al.*, 2014) permitted within organic production systems are used in combination with agro-management tools such as water and soil conservation.
- To increase organic matter within the soil, more sustainable crop production strategies in combination with minimum tillage are being utilised. To determine the most promising available novel cultural control strategies targeting the *Fusarium* inoculum within crop debris, *Brassica* biofumigation (Ngala *et al.*, 2015; Perniola *et al.*, 2014), as well as accelerated biodegradation (Hofgaard, 2012) with an extract or competitive fungi (Matarese *et al.*, 2012) are being explored.
- To increase the climate change resilience of EU-maize farmers and thus reduce susceptibility of maize to aflatoxin contamination, MyToolBox is testing two measures: (1) selecting appropriate genotypes of maize with various traits (e.g. drought resistance, early

mature plant cultivars) in field conditions that are likely to reduce aflatoxin contamination; and (2) applying atoxigenic *Aspergillus* strains that are out-competing toxigenic *Aspergillus* strains, which is a novel and proven technology developed in the USA and Africa (Atehnkeng *et al.*, 2008; Bandyopadhyay *et al.*, in press; Brown *et al.*, 1991; Probst *et al.*, 2011), adapted to Europe in collaboration with the international experts from those continents.

- To assist short term decision making regarding fungicide application, harvest timing, and long term planning, available forecasting models regarding DON contamination in wheat and maize are being validated, combined and extended to cover the most important wheat and maize growing areas in Europe. Previously developed weather-driven forecasting models were validated to a limited extent, mainly in the regions where they have been developed, and mostly to predict *Fusarium* head blight or DON contamination specifically in wheat. By synergising and extending existing models,

MyToolBox will therefore significantly improve the forecasting of potential DON contamination in major grain crops in Europe.

Post-harvest objectives

- To facilitate effective integrated control of moulds and mycotoxins and to provide an early warning of risks, and thus enable implementation of effective prevention/remedial actions, real-time post-harvest decision support systems for storage of cereals and peanuts are being developed. They are based on the sensitive indicator of CO₂, as well as temperature and relative humidity, and will be tested for local conditions in European and Chinese cereal and peanut silos.
- To achieve significant improvements in sorting of dried figs and to also improve the work environment dramatically, MyToolBox develops a non-invasive system for the real-time sorting of contaminated dried figs, based on visible and near infrared spectral imaging and appropriate algorithms. Traditionally, to remove units contaminated with aflatoxins, dried figs are sorted by hand (Benalia *et al.*, 2013; Oezlueymak, 2014; Patel *et al.*, 2012). These require working conditions in the dark, under UV light, which is potentially harmful to the health of the employees (Ichihashi *et al.*, 2003; Kammeyer and Luiten, 2015; Lavker *et al.*, 1995). Therefore, besides contributing to reduced contamination of mycotoxins in dried fig batches (with 5% or less probability of the batch average concentration of aflatoxin B₁ failing to meet the EU limits), MyToolBox will also contribute to occupational health improvements.
- Innovative pre-milling and milling strategies are combined and scaled-up, which will lead to more accurate separation of grain tissues with characterised (different) mycotoxin contamination levels. Milling strategies, such as debranning and micronisation, are efficient, but individual tissues of the grains are often inhomogeneous and not well confined. By-products, usually intended for animal feed but also exploitable for wholegrain food products, contain about 15% proteins, 40% starch and 40% fibre with levels of mycotoxins which can exceed legal limits (Delcour *et al.*, 2012). The synergistic potential of debranning, micronisation and dry- or wet-turbo separation will minimise the mycotoxin concentration in raw materials and the final wheat products destined for the consumer.
- Within MyToolBox, the fate of mycotoxins as well as their modified forms and co-occurrences in naturally contaminated raw materials (with simultaneous co-occurrence of different mycotoxins) will be monitored during thermal food processing at an industrial scale with Barilla (Parma, Italy). These measures are expected to provide (1) structural information of altered chemical structures (using newest LC-HRMS and metabolomics techniques (Bueschl *et al.*, 2014)); and (2) novel

bakery production protocols, controlling for instance fermentation, temperature/time and recipes, to develop a sound basis to establish process factors to reduce risks as part of a post-harvest intervention strategy (Bergamini *et al.*, 2010; Generotti *et al.*, 2015). These efforts cater to the need to obtain further data on the fate of masked/modified mycotoxins, as defined by EFSA in 2014 (EFSA CONTAM Panel, 2014), during food and feed processing. Hence, MyToolBox complements previous studies in this area by adding results of in-situ tests with practical applicability to lab-scale experiments that have used artificially contaminated material at high levels (Meca *et al.*, 2013; Valle-Algarra *et al.*, 2009; Voss and Snook, 2010).

- To study the use of mycotoxin contaminated batches in biofuel production MyToolBox partners Südzucker (Mannheim, Germany) and BIOMIN (Getzersdorf, Austria) are investigating the reduction of the mycotoxin content in the by-product DDGS, for which recently available novel enzymes will be utilised in technical scale fermentation processes. Maize-based DDGS can be contaminated with high levels and a broad spectrum of mycotoxins (US Grains Council, 2012; Rodrigues and Chin, 2011; Schaafsma *et al.*, 2009; Zachariasova *et al.*, 2014; Zhang and Caupert, 2012), resulting in losses in feed production. Micro-organisms can degrade mycotoxins but the efficacy for a safe use to efficiently produce biogas (Hanschmann, 2008) and bioethanol (Kłosowski *et al.*, 2010) by fermentation of mycotoxin-contaminated batches must still be determined. Therefore, the performance of specific mycotoxin-degrading microorganism strains (Vekiru *et al.*, 2010) and novel enzymes provided by partner BIOMIN, on the detoxification of DON (EC, 2013) and fumonisins (EC, 2014) during biogas and bioethanol production are being evaluated.

Objectives throughout the food and feed chain including dissemination

- To provide decision-support in mycotoxin management to different end-users along the food and feed chain, an integrated, dynamic, MyToolBox e-platform is being designed, tested and validated. This e-platform will provide users with information, tools and guidance concerning actions that can be taken to reduce fungal contamination, leading to reductions in contamination-levels of mycotoxins in the food and feed chains. The MyToolBox e-platform will be accessible to farmers, food processors and other actors along the chain through all mainstream interfaces (PCs, tablets or smartphones).
- To ensure economic feasibility, cost-effective sampling and monitoring strategies for mycotoxin management applied over the entire food and feed chain are adapted and integrated in the MyToolBox e-platform.
- To strengthen international cooperation in research and innovation for mycotoxins, all work packages reflect

collaborative actions and tasks between partners from two or more different countries. To invigorate the ties between China and Europe in the area of mycotoxin reduction, MyToolBox has initiated a formal EU-China dialogue through an EU-China Mycotoxin Forum.

3. Expected impact

Overall, the impacts of an integrated toolbox for mycotoxin management, as developed within MyToolBox shall be a reduced number of incidents and more transparent supply chains. The developed and applied pre- and post-harvest measures, including reliable monitoring of crops and proper risk assessment strategies, should lead to significant economic and food safety benefits and improvements in mycotoxin management along the food and feed chains. Moreover, MyToolBox shall contribute to improved consumer trust in imported food and feed. Members of EFSA will be on the advisory board to ensure European citizens are well informed regarding risks in the food and feed chain.

Building on the existing knowledge, MyToolBox integrates the wealth of knowledge on mitigation and reduction of mycotoxins in food and feed in one single online decision support platform, leading to more efficient use of resources. In particular unexpected contamination of grains can cause huge losses: For example, the long term costs for maize based feed contaminated with aflatoxins are estimated to cost Serbia hundreds of millions of Euros per year. Thus, it is expected that farmers and other stakeholders along the food and feed chain will benefit financially by saving resources (e.g. fertilisers, seeds, etc.) and by lowering the risk of rejection of their products due to unexpected contamination. As such, mycotoxin-related border rejections noted by RASFF are expected to decrease further (Anonymous, 2014).

Pre-harvest impacts

The pre-harvest impact of the MyToolBox work will be measured by comparing the achieved reduction in mycotoxin content and increase in grain yield, based on field trials, between the conventional agricultural systems to the newly designed systems. The use of biopesticides and biofumigation or debris treatment in EU field trials each should reduce the DON level in wheat by 50% (Paul *et al.*, 2008). However, the combined effect is yet to be explored. In EU field trials the aflatoxin content of resistant maize cultivars will be compared to susceptible cultivars, which is expected to be lower than previously achieved 70% (Windham and Williams, 1998). The aflatoxin contents of maize (EU) and peanuts (China) after treatment with atoxigenic fungal strains will be compared to non-treated crops in field trials. The aim is to achieve a 90% reduction in aflatoxin contamination (Dorner, 2004).

Post-harvest impacts

The post-harvest impact of the real-time sensors developed in MyToolBox will be measured by comparing the achieved reduction in mycotoxin content of cereals in trials in grain silos compared to conventional practices. This is expected to contribute to approximately 15% savings (~100 M€/year). While it is impossible to accurately determine the reduction in costs, it is estimated that costs for monitoring could be reduced by at least 30% at all stages of the food and feed supply chain. This impact will be measured by comparing the estimated current costs for monitoring by EU governments and industry to the newly designed monitoring systems. Overall, MyToolBox aims to reduce post-harvest losses, due to mould spoilage and mycotoxins, pests in food/feed chains, in the EU and its associated countries by at least 15%.

The novel non-invasive real-time sorting of dried figs is expected to result in maximum 5% probability of the batch average concentration of aflatoxin B₁ failing to meet the EU limits as compared to conventional sorting. The developed innovative milling and baking techniques aim to achieve a 50 and 15% mycotoxin reduction in the finished food products respectively. New information on the presence of mycotoxins and their modified forms resulting from food processing will be forwarded to EFSA as a relevant contribution for its overall risk assessment of mycotoxins in foods.

The effect of the use of enzymes on mycotoxin reduction during biofuel production is being assessed in trials with and without addition of the enzymes. The impact will be a significant reduction of the DON- and fumonisin-content in DDGS as opposed to conventional production. In 2013, about 5.2 million m³ of bioethanol were produced from grains in Europe (and 88.8 million m³ worldwide), resulting in a production of about 4.2 million tons of DDGS in Europe. It is estimated that the losses in feed production due to mycotoxin-contaminated DDGS to be in the range of 15-20 million Euro annually for the swine industry alone. Therefore, a reduction of losses is expected due to (1) significantly lowering mycotoxin levels in DDGS; (2) improved efficiency of the bioethanol production of 10-30%; and (3) increased market value (>10%) of DDGS of low mycotoxin content. The outcome of these MyToolBox studies could also positively contribute to the current food vs fuel debate, as mycotoxin contaminated batches could then be safely employed to produce biofuels on the one hand, and high quality DDGS on the other hand.

Impacts throughout the food and feed chain including dissemination

Finally, the web-based MyToolBox platform will represent a one-stop-shop consisting of information and advisory modules, management tools, as well as technologies that

equip farmers and all other actors in the food and feed supply chain with systematic, pro-active, cost effective and affordable approaches for effective monitoring and reduction of the incidences of mycotoxin contamination in crops, food and feed. In addition, it will contribute to the reduction of stakeholders' dependence on expensive and wasteful end-point testing and segregation.

The EU-China partnership within MyToolBox is aimed to contribute to the standard settings for authorisation of mycotoxin-detoxifying feed additives in China, whereby current EU guidelines for the registration of detoxifying feed additives shall serve as examples for possible adaptations of Chinese legislation. This is important as currently about 475 million pigs are farmed in China, representing approximately half of the world's total pig population. Since up to 24% of complete feed for pigs have exceeded the maximum limits set for DON levels by the Chinese authorities (Li *et al.*, 2014). The need for more than 54,000 metric tons of detoxifying agents can be estimated in China, thus contributing to more efficient pork-meat production in China.

4. Conclusions

The MyToolBox project does not only pursue a field-to-fork approach to reduce mycotoxins in wheat, oats, maize, peanuts and dried figs, but also considers safe use options of mycotoxin contaminated batches such as microbial energy conversion for efficient production of biogas, bioethanol and safe DDGS. Up to now, different agricultural strategies including crop rotation, tillage practices, fungicide application and the use of resistant plant cultivars have been used to reduce the impact of fungal infection of crops and subsequent mycotoxin formation. As none of these strategies by themselves have been able to efficiently reduce the impact of fungal infection, a novel integrated approach is pursued, which also takes into account issues related to climate change. Consequently, MyToolBox combines a series of integrated pre- and post-harvest measures, which could enable a 20-90% reduction (depending on the type of commodity and intervention) in losses of crops due to fungal and mycotoxin contamination.

Acknowledgements

The authors are grateful to all the partners' contributions, in particular Marc Lemmens from the University of Natural Resources and Life Sciences, Vienna, Kurt Brunner from the Vienna University of Technology (current affiliation: Romer Labs), Hamide Senyuva from FoodLife International, Angel Medina Vaya from Cranfield University, Edurne Gaston Estanga from IRIS, Dragana Krajcinovic from the University of Novi Sad, Sara Elisabetta Legler from HORTA, Carsten Fauhl-Hassek from the German Bundesministerium für Risikobewertung, Habil Kalkan from Suleyman Demirel University, Monique Mourits

from Wageningen University, Ingerd Hofgaard from the Norwegian Institute of Bioeconomy Research, Liu Yang from the Chinese Academy of Agricultural Sciences – Institute of Food Science and Technology, Gerd Schatzmayr and Veronika Nagl from BIOMIN Holding GmbH, Songxue Wang from the Chinese Academy of State Administration of Grain, Michael Klingenberg from Suedzucker, Ruben Torregrosa from Axeb Biotech SL, Giel Nussgelder from Agro L V Limited, Nil Yetgin from Taris Figs Agricultural Sales Cooperatives Unions, Helena Stanko from D.O.O. Agrocentrum Becej, and Jinqian Wang from the Chinese Academy of Agricultural Sciences – Feed Research Institute. Finally, we would also like to thank the MyToolBox Project Officer Alessio Bacchielli for his guidance during the preparation of the grant agreement and the project's kick-off. This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 678012.

References

- Anonymous, 2013. EU: feed contaminated with aflatoxin. All About Feed, 11 March 2013 News. Available at: <http://tinyurl.com/z784mfh>.
- Anonymous, 2014. Decrease in EU mycotoxin alerts. All About Feed, 30 July 2014 News. Available at: <http://tinyurl.com/jgvjys9>.
- Atehnkeng, J., Ojiambo, P.S., Ikotun, T., Sikora, I.A., Cotty, P.J. and Bandyopadhyay, R., 2008. Evaluation of atoxigenic isolates of *Aspergillus flavus* as potential biocontrol agents for aflatoxin in maize. Food Additives and Contaminants Part A 25: 1264-1271.
- Bandyopadhyay, R., Ortega-Beltran, A., Akande, A., Mutegi, C., Atehnkeng, J., Kaptoge, L., Senghor, A.L., Adhikari, B.N. and Cotty, P.J., in press. Biological control of aflatoxins in Africa: current status and potential challenges in the face of climate change. World Mycotoxin Journal, DOI: <https://doi.org/10.3920/WMJ2016.2130>.
- Battilani, P., Toscano, P., Van der Fels-Klerx, H.J., Moretti, A., Camardo Leggieri, M., Brera, C., Rortais, A., Goumperis, T. and Robinson, T., 2016. Aflatoxin B₁ contamination in maize in Europe increases due to climate change. Scientific Reports 6: 24328. DOI: <http://dx.doi.org/10.1038/srep24328>.
- Benalia, S., Zimbalatti, G., Cubero, S., Prats, J.M., Bernardi, B., Chanona-Pérez, J.J., Blasco, J., 2013. Automatic sorting of dried figs based on computer vision systems. In: VII Congreso Ibérico de Agroingeniería y Ciencias Hortícolas, 26-29 August 2013, Madrid, Spain, article C-0533.
- Bergamini, E., Catellani, D., Dall'asta, C., Galaverna, G., Dossena, A., Marchelli, R., Suman, M., 2010. Fate of *Fusarium* mycotoxins in the cereal product supply chain: the deoxynivalenol (DON) case within industrial bread-making technology. Food Additives and Contaminants 27: 677-687.
- Bueschl, C., Kluger, B., Lemmens, M., Adam, G., Wiesenberger, G., Maschietto, V., Marocco, V., Strauss, J., Bödi, S., Thallinger, G.G., Krska, R. and Schuhmacher, R., 2014. A novel stable isotope labelling assisted workflow for improved untargeted LC-HRMS based metabolomics research. Metabolomics 10: 754-769.

- Brown, R., Cotty, P.J. and Cleveland, T.E., 1991. Reduction in aflatoxin content of maize by atoxigenic strains of *Aspergillus flavus*. *Journal of Food Protection* 54: 623-626.
- Delcour, J.A., Rouau, X., Courtin, C.M., Poutanen, K. and Ranieri, R., 2012. Technologies for enhanced exploitation of the health-promoting potential of cereals. *Trends in Food Science and Technology* 20: 1-9.
- Dorner, J.W., 2004. Biological control of aflatoxin contamination in crops. *Journal of Toxicology: Toxin Reviews* 23: 425-450.
- European Food Safety Authority (EFSA) CONTAM Panel, 2014. Opinion of the Scientific Panel on Contaminants in the Food Chain on a request from the Commission related to the presence of modified forms of certain mycotoxins in food and feed. *EFSA Journal* 12: 3916.
- European Commission (EC), 2013. Commission Implementing Regulation (EU) No 1016/2013 of 23 October 2013 concerning the authorisation of a preparation of a micro-organism strain DSM 11798 of the *Coriobacteriaceae* family as a feed additive for pigs. *Official Journal of the European Union L* 282: 36.
- European Commission (EC), 2014. Commission Implementing Regulation (EU) No 1115/2014 of 21 October 2014 concerning the authorisation of a preparation of fumonisin esterase produced by *Komagataella pastoris* (DSM 26643) as a feed additive for pigs. *Official Journal of the European Union L* 302: 51.
- European Commission (EC), 2015. European Commission Decision C (2015) 2453: Horizon 2020 Work Programme 2014-2015: Food security, sustainable agriculture and forestry, marine and maritime and inland water research and bioeconomy (revised). Available at: <http://tinyurl.com/hdqqnq8>.
- European Commission (EC), 2016. The Rapid Alert System for Food and Feed 2015 Annual Report. Available at: <http://tinyurl.com/h7l6v9x>.
- Eurostat, 2015a. Selling prices of crop products (absolute prices) – annual price from 2000 onwards). Available at: <http://tinyurl.com/z2sxbq6>.
- Eurostat, 2015b. Crops product – annual data. Available at: <http://ec.europa.eu/eurostat/data/database>.
- Forrer, H.R., Musa, T., Schwab, F., Jenny, E., Bucheli, T.D., Wettstein, F.E. and Vogelgsang, S., 2014. *Fusarium* Head Blight control and prevention of mycotoxin contamination in wheat with botanicals and tannic acid. *Toxins* 6: 830-849.
- Generotti, S., Cirlini, M., Malachova, A., Sulyok, M., Berthiller, F., Dall'Asta, C. and Suman, M., 2015. Deoxynivalenol & deoxynivalenol-3-glucoside mitigation through bakery production strategies: effective experimental design within industrial rusk-making technology. *Toxins* 7: 2773-2790.
- Hanschmann, G., 2008. Versuch zum Verhalten von DON in einer Biogasanlage. Sächsische Landesanstalt für Landwirtschaft, Fachbereich Landwirtschaftliches Untersuchungswesen. Available at: <http://tinyurl.com/gtcg8km>.
- Hofgaard, I.S., 2012. Reduction in the inoculum potential of *F. graminearum* on residues after treatment with Biowish crop or Trifender after sowing in spring. Unpublished internal report, Norwegian Institute for Agricultural and Environmental Research, Dep. of Fungal Plant Pathology, Ås, Norway.
- Ichihashi, M., Ueda, M., Budiyanto, A., Bito, T., Oka, M., Fukunaga, M., Tsuru, K. and Horikawa, T., 2003. UV-induced skin damage. *Toxicology* 189: 21-39.
- Jakšić, S., Živkov-Baloš, M., Prica, N., Mašić, Z., Nešić, K., Jajić, I. and Abramović, B., 2015. The influence of climatic factors in Serbia on mycotoxin production. In: Scientific Veterinary Institute 'Novi Sad': Proceedings of the First International Symposium of Veterinary Medicine (ISVM2015) 'One Health – New Challenges', Novi Sad, Serbia, pp. 166-172.
- Jovanović, M., Trailović, D., Kukulj, V., Nesic, S., Marinkovic, D., Nedeljkovic-Trailovic, J., Jakovac Strajn, B. and Milicevic, D., 2015. An outbreak of fumonisin toxicosis in horses in Serbia. *World Mycotoxin Journal* 8: 387-391.
- Kammeyer, A. and Luiten, R.M., 2015. Oxidation events and skin aging. *Ageing Research Reviews* 21: 16-29.
- Kłosowski, G., Mikulski, D., Grajewski, J. and Błajet-Kosicka, A., 2010. The influence of raw material contamination with mycotoxins on alcoholic fermentation indicators. *Bioresource Technology* 101: 3174-3152.
- Lavker, R.M., Gerberick, G.F., Veres, D., Irwin, C.J. and Kaidbey, K.H., 1995. Cumulative effects from repeated exposures to suberythemal doses of UVB and UVA in human skin. *Journal of the American Academy of Dermatology* 32: 53-62.
- Li, X.Y., Zhao, L., Fan, Y., Jia, Y., Sun, L., Ma, S., Ji, C., Ma, Q. and Zhang, J., 2014. Occurrence of mycotoxins in feed ingredients and complete feeds obtained from the Beijing region of China. *Journal of Animal Science and Biotechnology* 5: 37-44.
- Lubulwa, G.A.S., Siriacha, P., Markwell, P.J. and Pitt, J.I., 2015. Estimating the burden of market loss due to aflatoxins in maize: methods and estimates for Thailand. *World Mycotoxin Journal* 8: 459-464.
- Matacic, C., 2016. Fungal toxins are poisoning Africa's children, says new report. *Science, News* 18 February 2016. Available at: <http://tinyurl.com/jphng2f>.
- Matarese, F., Sarrocco, S., Gruber, S., Seidl-Seiboth, V. and Vannacci, G., 2012. Biocontrol of *Fusarium* head blight: interactions between *Trichoderma* and mycotoxigenic *Fusarium*. *Microbiology* 158: 98-106.
- Meca, G., Zhou, T., Li, X.-Z. and Manes, J., 2013. Beauvericin degradation during bread and beer making. *Food Control* 34: 1-8.
- Nährer, K. and Kovalsky, P., 2014. The Biomin mycotoxin survey: identifying the threats in 2013. *Science and Solutions*, April 2014. Available at: <http://tinyurl.com/ztnjsas>.
- Ngala, B.M., Haydock, P.P., Woods, S. and Back, M.A., 2015. Biofumigation with *Brassica juncea*, *Raphanus sativus* and *Eruca sativa* for the management of field populations of the potato cyst nematode *Globodera pallida*. *Pest Management Science* 71: 759-769.
- Norwegian Scientific Committee for Food Safety (VKM), 2013. Risk assessment of mycotoxins in cereal grain in Norway. Opinion of the Scientific Steering Committee of the Norwegian Scientific Committee for Food Safety. No. 10/004-04-Final. Available at: <http://tinyurl.com/jkghpqx>.
- Oezlueoymak, O.B., 2014. Development of a UV-based imaging system for real-time detection and separation of dried figs contaminated with aflatoxins. *Journal of Agricultural Sciences* 20: 302-316.

- Patel, K.K., Kar, A., Jha, S.N. and Khan, M.A., 2012. Machine vision system: a tool for quality inspection of food and agricultural products. *Journal of Food Science and Technology* 49: 123-141.
- Paul, P.A., Lipps, P.E., Hershman, D.E., McMullen, M.P., Draper, M.A. and Madden, L.V., 2008. Efficacy of triazole-based fungicides for *Fusarium* Head Blight and deoxynivalenol control in wheat: a multivariate meta-analysis. *Phytopathology* 98: 999-1011.
- Perniola, O.S., Staltari, S., Chorzempa, S.E., Astiz-Gassó, M.M. and Del Carmen Molina, M., 2014. Biological control of *Fusarium graminearum*: use of *Trichoderma* spp. and biofumigation with aerial part of *Brassica juncea*. *Revista De La Facultad De Ciencias Agrarias* 46: 45-56.
- Probst, C., Bandyopadhyay, R., Price, L.E. and Cotty, P.J., 2011. Identification of atoxigenic *Aspergillus flavus* isolates to reduce aflatoxin contamination of maize in Kenya. *Plant Disease* 95: 212-218.
- Rodrigues, I. and Chin, L.J., 2012. A comprehensive survey on the occurrence of mycotoxins in maize dried distillers' grain and solubles sourced worldwide. *World Mycotoxin Journal* 5: 83-88.
- Schaafsma, A.W., Limay-Rios, V., Paul, D.E. and Miller, J.D., 2009. Mycotoxins in fuel ethanol co-products derived from maize: a mass balance for deoxynivalenol. *Journal of the Science of Food and Agriculture* 89: 157-1580.
- Schmale, D.G. and Munkvold, G.P., 2009. Mycotoxins in crops: a threat to human and domestic animal health. American Phytopathological Society, St. Paul, MN, USA. Available at: <http://tinyurl.com/hjxnz86>.
- US Grains Council, 2012. A guide to distiller's dried grains with solubles (DDGS). 3rd ed. Available at: <http://tinyurl.com/j6fwvb9>.
- Valle-Algarra, F.M., Mateo, E.M., Medina, A., Mateo, R., Gimeno-Adelantado, J.V. and Jimenez, M., 2009. Changes in ochratoxin A and type A and type B trichothecenes contained in wheat flour during dough fermentation and bread making. *Food Additives and Contaminants Part A* 26: 896-906.
- Vardon, P.J., McLaughlin, C. and Nardinelli, C., 2003. Potential economic costs of mycotoxins in the United States. In: CAST report: Mycotoxins; risks in plant, animal and human systems. Task Force Report no. 139, CAST, Ames, Iowa, USA, pp. 136-142.
- Vekiru, E., Hametner, C., Mitterbauer, R., Rechthaler, J., Adam, G., Schatzmayr, G., Krska, R. and Schuhmacher, R., 2010. Cleavage of zearalenone by *Trichosporon mycotoxinivorans* to a novel nonestrogenic metabolite. *Applied and Environmental Microbiology* 76: 2353-2359.
- Voss, K.A. and Snook, M.E., 2010. Stability of the mycotoxin deoxynivalenol (DON) during the production of flour-based foods and wheat flake cereal. *Food Additives and Contaminants Part A* 27: 1694-1700.
- Wild, C.P., Miller, J.D. and Groopman, J.D., 2015. Mycotoxin control in low- and middle-income countries. Working Group Report No. 9, International Agency for Research on Cancer, Lyon, France.
- Windham, G.L. and Williams, W.P., 1998. *Aspergillus flavus* infection and aflatoxin accumulation in resistant and susceptible maize hybrids. *Plant Disease* 82: 281-284.
- Wu, F. and Munkvold, G.P., 2008. Mycotoxins in ethanol co-products: modeling economic impacts on the livestock industry and management strategies. *Journal of Agricultural and Food Chemistry* 56: 3900-3911.
- Zachariasova, M., Dzuman, Z., Veprikova, Z., Hajkova, K., Jiru, M., Vaclavikova, M., Pospichalova, M., Florian, M. and Hajslova, J., 2014. Occurrence of multiple mycotoxins in European feedingstuffs, assessment of dietary intake by farm animals. *Animal Feed Science and Technology* 193: 124-140.
- Zhang, Y. and Caupert, J., 2012. Survey of mycotoxins in U.S. distiller's dried grains with solubles from 2009 to 2011. *Journal of Agricultural and Food Chemistry* 60: 539-543.

